


Vali-e-Asr University
of Rafsanjan

Wavelets and Linear Algebra

<http://wala.vru.ac.ir>



Pseudoframe multiresolution structure on locally compact abelian groups

Hamide Azarmi^a, Rajab Ali Kamyabi Gol^a, Mohammad Janfada^{a,*}

^aDepartment of pure Mathematics, Ferdowsi University of Mashhad, P.O. Box 1159 Mashhad 91775, Islamic Republic of Iran.

ARTICLE INFO

Article history:

Received 31 December 2015

Accepted 10 November 2016

Available online 25 December 2016

Communicated by Abdolaziz Abdollahi

Keywords:

Pseudoframe,
generalized
multiresolution
structure.

2000 MSC:

42C15, 43A70

ABSTRACT

Let G be a locally compact abelian group. The concept of generalized multiresolution structure (GMS) in $L^2(G)$ is discussed which is a generalization of GMS in $L^2(\mathbb{R})$. Basically, a GMS in $L^2(G)$ consists of an increasing sequence of closed subspaces of $L^2(G)$ and a pseudoframe of translation type at each level. Also, the construction of affine pseudoframes for $L^2(G)$ based on a GMS is presented.

© (2016) Wavelets and Linear Algebra

*Corresponding author

Email addresses: azarmi_1347@yahoo.com (Hamide Azarmi), kamyabi@um.ac.ir (Rajab Ali Kamyabi Gol), janfada@um.ac.ir (Mohammad Janfada)

1. Introduction and preliminary

In 1946, Gabor introduced an interesting approach to signal decomposition [12]. In 1952, Duffin and Schaeffer abstracted Gabor’s method to introduce the notion of frame in nonharmonic Fourier analysis [9]. The idea of Duffin and Schaeffer was not continued until 1986 when Daubechies et al. in [8], applied the theory of frames to wavelets and Gabor transforms. After their work, the theory of frames began to be studied widely and deeply by many authors (see [3]-[5], for example). Today, the theory of frames has been applied to signal processing, image processing, data compressing and sampling theory and so on.

A sequence $\{x_n\}_{n \in \mathbb{Z}}$ in a separable Hilbert space \mathcal{H} is called a frame for \mathcal{H} if there exist constants $A, B > 0$ such that

$$A\|x\|^2 \leq \sum_n |\langle x, x_n \rangle|^2 \leq B\|x\|^2, \quad (x \in \mathcal{H}). \tag{1.1}$$

If the right inequality holds, then $\{x_n\}_{n \in \mathbb{Z}}$ is said to be a Bessel sequence. It is well known that for any frame $\{x_n\}_{n \in \mathbb{Z}}$ there exists another frame $\{x_n^*\}_{n \in \mathbb{Z}}$ in \mathcal{H} , namely dual frame of $\{x_n\}_{n \in \mathbb{Z}}$, such that for any $x \in \mathcal{H}$

$$x = \sum_n \langle x, x_n^* \rangle x_n = \sum_n \langle x, x_n \rangle x_n^*.$$

The concept of a multiresolution analysis (MRA) was first introduced by Mallat [21] and Meyer [22]. It is a general framework for constructing orthonormal wavelet bases for $L^2(\mathbb{R})$ of the form $\{2^{j/2}\psi(2^j \cdot - k)\}_{j,k \in \mathbb{Z}}$. The MRA-based compactly supported orthonormal wavelet systems were constructed by Daubechies [7].

Frame multiresolution analysis (FMRA) as a generalization of MRA, introduced by Benedetto and Li in [2].

As usual, we define the following operators on $L^2(\mathbb{R})$ by

$$(\tau_b f)(x) = f(x - b), \quad (Df)(x) = 2^{1/2} f(2x).$$

The parameter b in the first operator can be an arbitrary real number. A frame multiresolution analysis (FMRA) for $L^2(\mathbb{R})$ consists of a sequence of closed linear subspaces $\{V_j\}_{j \in \mathbb{Z}}$ of $L^2(\mathbb{R})$ and a function $\phi \in V_0$ such that

1. $V_j \subseteq V_{j+1}$,
2. $\overline{\cup_j V_j} = L^2(\mathbb{R}), \cap_j V_j = \{0\}$,
3. $V_j = D^j V_0$,
4. $f \in V_0$ implies that $\tau_k f \in V_0$, for all $k \in \mathbb{Z}$,
5. $\{\tau_k \phi : k \in \mathbb{Z}\}$ is a frame for V_0 .

In the above definition, if $\{\tau_k \phi : k \in \mathbb{Z}\}$ is an orthonormal basis for V_0 , then $\{V_j, \phi\}_{j \in \mathbb{Z}}$ forms a multiresolution analysis (MRA) for $L^2(\mathbb{R})$. For more details on FMRA, one can see [1], [4].

Dahlke in [6], generalized the notion of MRA to locally compact abelian groups and proved an existence theorem based on generalized B-splines. For some groups G different from \mathbb{R}^d , multiresolution analysis were studied in [6], [10], [15] and [16]. In particular, the group analogues

of the B-spline wavelet bases in $L^2(\mathbb{R})$ are defined in [6] and [10]. In [14], conditions under which a function generates a multiresolution analysis on a locally compact abelian groups were investigated.

The notion of generalized multiresolution structure (GMS) in $L^2(\mathbb{R})$ was introduced in [18]. Basically, the GMS consists of an increasing sequence of closed subspace of $L^2(\mathbb{R})$, with a pseudoframe of translates at each level. Let $\{\tau_k\phi\}_{k \in \mathbb{Z}}$ and $\{\tau_k\phi^*\}_{k \in \mathbb{Z}}$ be two sequences in $L^2(\mathbb{R})$ and \mathcal{X} be a closed subspace of $L^2(\mathbb{R})$. We say $\{\tau_k\phi\}_{k \in \mathbb{Z}}$ forms a pseudoframe of translates for \mathcal{X} with respect to $\{\tau_k\phi^*\}_{k \in \mathbb{Z}}$ if

$$x = \sum_k \langle x, \tau_k\phi^* \rangle \tau_k\phi, \quad (x \in \mathcal{X}).$$

In a more general case, let \mathcal{X} be a closed subspace of a separable Hilbert space \mathcal{H} . Let $\{x_n\}_{n \in \mathbb{Z}} \subset \mathcal{H}$ be a Bessel sequence with respect to \mathcal{X} , and let $\{x_n^*\}_{n \in \mathbb{Z}} \subset \mathcal{H}$ be a Bessel sequence in \mathcal{H} . We say $\{x_n\}_{n \in \mathbb{Z}}$ is a pseudoframe for the subspace \mathcal{X} (PFFS) with respect to $\{x_n^*\}_{n \in \mathbb{Z}}$ if

$$x = \sum_k \langle x, x_k \rangle x_k^*, \quad (x \in \mathcal{X}).$$

$\{x_n^*\}_{n \in \mathbb{Z}}$ is called a dual pseudoframe (or PFFS-dual) of $\{x_n\}_{n \in \mathbb{Z}}$ for the subspace \mathcal{X} , (see [19] and [20] for more details).

For the reader's convenience, we report a number of definitions. In this paper, we assume that G is a locally compact abelian group and Γ is a uniform lattice in G , that is Γ is a discrete subgroup that $\frac{G}{\Gamma}$ is compact. If Γ is a uniform lattice, then Γ^\perp defined by $\{\xi \in \widehat{G} : \xi(\Gamma) = 1\}$ is a uniform lattice in \widehat{G} , where \widehat{G} is the dual group of G [13].

Let $\pi : \Gamma \rightarrow U(L^2(G))$ be the translation representation which is defined by $(\pi_\gamma f)(x) = f(x\gamma^{-1})$. Let σ be a (continuous) unitary operator on $L^2(G)$ with the property $\sigma^{-1}\pi_\gamma\sigma = \pi_{\alpha(\gamma)}$, where α is an injective endomorphism on Γ . Also let δ be a (continuous) topological automorphism on G such that $\delta(\Gamma) \subset \Gamma$.

The Fourier transform $\widehat{\cdot} : L^1(G) \rightarrow C_0(\widehat{G})$, $f \mapsto \widehat{f}$, is defined by $\widehat{f}(\xi) = \int_G f(x)\overline{\xi(x)}dx$. The Fourier transform can be extended to a unitary isomorphism from $L^2(G)$ to $L^2(\widehat{G})$, known as the Plancherel transform (see [11]).

In this paper, we generalize the notion of GMS for $L^2(G)$. In Section 2, a necessary and sufficient condition for existence of pseudoframes for Paley-Wiener subspaces is studied. Based on this concept, a formal definition of a GMS for a locally compact abelian group is given in Section 3. Consequently, construction methods for GMSs are also explained. Furthermore, a construction that allows us to obtain affine pseudoframes associated with such a GMS is given in Section 4.

Our conclusions are mainly generalizations of results of Li in [18].

2. Existence of pseudoframes for subspaces of $L^2(G)$

There are some examples in $L^2(G)$ such that $\{\pi_\gamma\phi\}_{\gamma \in \Gamma}$ is neither generates MRA nor FMRA. Nevertheless a stable expansion for elements of a closed subspace of $L^2(G)$ exists in term of $\{\pi_\gamma\phi\}_{\gamma \in \Gamma}$ (see [1, 17, 19, 20]). The following is such an example.

Example 2.1. Let $G = \mathbb{R} \times \mathbb{R}$ and $\Gamma = \mathbb{Z} \times \mathbb{Z}$. Consequently, $\hat{\Gamma} = \hat{\mathbb{Z}} \times \hat{\mathbb{Z}} = \mathbb{T} \times \mathbb{T}$. Define a function $\phi \in L^2(G)$ such that

$$\hat{\phi}(\gamma_1, \gamma_2) = \begin{cases} 1 & (\gamma_1, \gamma_2) \in [-\frac{1}{4}, \frac{1}{4}]^2 \\ \text{decaying to zero continuously} & (\gamma_1, \gamma_2) \in [-\frac{1}{2}, \frac{1}{2}]^2 - [-\frac{1}{4}, \frac{1}{4}]^2 \\ 0 & \text{outside of } [-\frac{1}{2}, \frac{1}{2}]^2 \end{cases}$$

Let $\Delta := [-\frac{1}{4}, \frac{1}{4}]^2$ and define $V_0 := PW_\Delta$ that is the Paley-Wiener space, the space of all band-limited functions with bandwidth in Δ (see [1]). By Shannon sampling theorem for $L^2(\mathbb{R}^d)$ [23], and for all $f \in PW_\Delta$ we have

$$f(x, y) = \sum_{(m,n) \in \mathbb{Z} \times \mathbb{Z}} f(m, n) \pi_{(m,n)} \phi(x, y).$$

Since the function $\Phi(\gamma_1, \gamma_2) = \sum_{(m,n) \in \mathbb{Z} \times \mathbb{Z}} |\hat{\phi}(\gamma_1 + m, \gamma_2 + n)|^2$ is continuous, we are not able to find the lower frame bound for $\{\pi_{(m,n)} \phi\}_{(m,n) \in \mathbb{Z} \times \mathbb{Z}}$. So $\{\pi_{(m,n)} \phi\}_{(m,n) \in \mathbb{Z} \times \mathbb{Z}}$ cannot be a frame for the closure of the span of $\{\pi_{(m,n)} \phi(t_1, t_2) : (m, n) \in \mathbb{Z} \times \mathbb{Z}\}$.

Also, from the fact that $\phi \notin PW_\Delta$, the sequence $\{\pi_{(m,n)} \phi\}_{(m,n) \in \mathbb{Z} \times \mathbb{Z}}$ is not a frame for PW_Δ . Whereas, if we define $V_j := PW_{2^j \Delta}$, $j \in \mathbb{Z}$, we have $V_j \subseteq V_{j+1}$, $\bigcup_j V_j = L^2(G)$ and $\bigcap_j V_j = \{0\}$.

Example 2.1 leads us to define the concept of pseudoframes on $L^2(G)$ for a locally compact abelian group G . For a uniform lattice Γ in G , let $\pi : \Gamma \rightarrow U(L^2(G))$ be the translation representation which is defined by $(\pi_\gamma f)(x) = f(x\gamma^{-1})$.

Definition 2.2. Let $\phi, \psi \in L^2(G)$ and \mathcal{X} be a closed subspaces of $L^2(G)$. The family $\{\pi_\gamma \phi\}_{\gamma \in \Gamma}$ is said to be a pseudoframe with respect to $\{\pi_\gamma \psi\}_{\gamma \in \Gamma}$ for \mathcal{X} , if for every $x \in \mathcal{X}$,

$$x = \sum_{\gamma \in \Gamma} \langle x, \pi_\gamma \psi \rangle \pi_\gamma \phi.$$

It is important to note that $\pi_\gamma \phi$ and $\pi_\gamma \psi$ need not be contained in \mathcal{X} . Also they are not generally commutable, this means there exists $x \in \mathcal{X}$ such that the following is not true,

$$x = \sum_{\gamma \in \Gamma} \langle x, \pi_\gamma \phi \rangle \pi_\gamma \psi.$$

In the following theorem, we are going to find a sufficient and necessary condition for the functions ϕ and ψ such that their translations forms a pseudoframe.

Theorem 2.3. Let $\phi \in L^2(G)$ be such that $|\hat{\phi}| > 0$, a.e. and $\hat{\phi}$ be zero outside of $\hat{\Gamma}$. For a fixed $c > 0$, let $\Delta := \{\gamma \in \hat{G} : |\hat{\phi}(\gamma)| \geq c\}$ be closed and let $V_0 := \{f \in L^2(G) : \text{supp } \hat{f} \subseteq \Delta\}$. For a $\psi \in L^2(G)$, $\{\pi_\gamma \phi\}_{\gamma \in \Gamma}$ forms a pseudoframe for V_0 with respect to $\{\pi_\gamma \psi\}_{\gamma \in \Gamma}$ if and only if

$$\widehat{\phi \bar{\psi}} \chi_\Delta = \chi_\Delta, \quad \text{a.e.}$$

Moreover, if ψ satisfies $|\hat{\psi}| > 0$ on $\hat{\Gamma}$, and the above equality holds, then $\pi_\gamma \phi$ and $\pi_\gamma \psi$ commute, in the sense that for any $x \in \mathcal{X}$,

$$x = \sum_{\gamma \in \Gamma} \langle x, \pi_\gamma \psi \rangle \pi_\gamma \phi = \sum_{\gamma \in \Gamma} \langle x, \pi_\gamma \phi \rangle \pi_\gamma \psi.$$

Proof. If $f \in V_0$, then $supp \hat{f} \subseteq \Delta$. By the assumptions, $supp \hat{\phi} \subseteq \hat{\Gamma}$. Now compactness of $\hat{\Gamma}$ implies that $supp \hat{\phi}$ is compact. On the other hand Δ is closed and so the fact that $\Delta \subseteq supp \hat{\phi}$ implies that Δ is compact. Hence $supp \hat{f}$ is compact.

By Weyl's formula we have

$$\begin{aligned} \left(\sum_{\eta \in \Gamma} \langle f, \pi_\eta \psi \rangle \pi_\eta \phi\right)(\gamma) &= \sum_{\eta \in \Gamma} \langle f, \pi_\eta \psi \rangle \hat{\phi}(\gamma) \overline{\gamma(\eta)} \\ &= \sum_{\eta \in \Gamma} \langle \hat{f}, \pi_\eta \hat{\psi} \rangle \hat{\phi}(\gamma) \overline{\gamma(\eta)} \\ &= \sum_{\eta \in \Gamma} \int_{\hat{G}} \hat{f}(\lambda) \overline{\hat{\psi}(\lambda)} \lambda(\eta) d\lambda \hat{\phi}(\gamma) \overline{\gamma(\eta)} \\ &= \sum_{\eta \in \Gamma} \int_{\hat{G}/\Gamma^\perp} \sum_{\xi \in \Gamma^\perp} \hat{f}(\xi\lambda) \overline{\hat{\psi}(\xi\lambda)}(\xi\lambda)(\eta) d\mu(\lambda\Gamma^\perp) \hat{\phi}(\gamma) \overline{\gamma(\eta)} \\ &= \hat{\phi}(\gamma) \sum_{\eta \in \Gamma} \left(\sum_{\xi \in \Gamma^\perp} \hat{f}(\xi\lambda) \overline{\hat{\psi}(\xi\lambda)}\right)^\vee(\eta) \overline{\gamma(\eta)} \\ &= \hat{\phi}(\gamma) \left(\sum_{\xi \in \Gamma^\perp} \hat{f}(\xi\lambda) \overline{\hat{\psi}(\xi\lambda)}\right)^\vee(\gamma) \\ &= \hat{\phi}(\gamma) \sum_{\xi \in \Gamma^\perp} \hat{f}(\xi\gamma) \overline{\hat{\psi}(\xi\gamma)}. \end{aligned}$$

The facts that $\hat{\phi}$ and \hat{f} are zero outside of $\hat{\Gamma}$ imply that the only nonzero term in the last summation is $\hat{\phi}(\gamma) \hat{f}(\gamma) \overline{\hat{\psi}(\gamma)}$. So

$$\hat{\phi} \overline{\hat{\psi}} \chi_\Delta = \chi_\Delta, \quad a.e.$$

For two Bessel families $\{\pi_\gamma \phi\}_{\gamma \in \Gamma}$ and $\{\pi_\gamma \psi\}_{\gamma \in \Gamma}$ in $L^2(G)$, define $U, V : L^2(G) \rightarrow l^2(\Gamma)$ by $U(f) = \{\langle f, \pi_\gamma \phi \rangle\}_{\gamma \in \Gamma}$ and $V(f) = \{\langle f, \pi_\gamma \psi \rangle\}_{\gamma \in \Gamma}$. From [19], we know that $\{\pi_\gamma \phi\}_{\gamma \in \Gamma}$ forms a pseudoframe with respect to $\{\pi_\gamma \psi\}$ for X if and only if

$$V^* U P = P,$$

where P is the orthogonal projection on X and also $\{\pi_\gamma \phi\}_{\gamma \in \Gamma}$ and $\{\pi_\gamma \psi\}_{\gamma \in \Gamma}$ commute if and only if $V^* U P = P = P U^* V$, where V^* and U^* are the adjoints of U and V , respectively. Indeed, we have

$$\begin{aligned} \langle f, P U^* V g \rangle &= \langle P f, U^* V g \rangle \\ &= \overline{\langle U^* V g, P f \rangle} \\ &= \sum_{\gamma \in \Gamma} \overline{\langle g, \pi_\gamma \psi \rangle \langle \pi_\gamma \phi, P f \rangle} \\ &= \sum_{\gamma \in \Gamma} \langle P f, \pi_\gamma \phi \rangle \langle \pi_\gamma \psi, g \rangle \\ &= \langle P f, \sum_{\gamma \in \Gamma} \langle g, \pi_\gamma \psi \rangle \pi_\gamma \phi \rangle \\ &= \langle P f, g \rangle = \langle f, P g \rangle \end{aligned}$$

which implies that $PU^*V = P$. Thus $\{\pi_\gamma\phi\}_{\gamma\in\Gamma}$ and $\{\pi_\gamma\psi\}_{\gamma\in\Gamma}$ commute. □

The following result of Li (Theorem 1, [18]) is a consequence of Theorem 2.3.

Corollary 2.4. *Let $\phi \in L^2(\mathbb{R})$ be such that $|\hat{\phi}| > 0$ a.e. on a connected neighborhood of 0 in $[-\frac{1}{2}, \frac{1}{2})$ and $|\hat{\phi}| = 0$, a.e. otherwise. Define $\Omega = \{\gamma \in \hat{\mathbb{R}} : |\hat{\phi}| \geq c > 0\}$, and let $V_0 := PW_\Omega = \{f \in L^2(\mathbb{R}) : \text{supp}(\hat{f}) \subseteq \Omega\}$. Then, for a $\psi \in L^2(\mathbb{R})$ and $\{\pi_k\phi\}_{k\in\mathbb{Z}}$ is a pseudoframe of translates for V_0 with respect to $\{\pi_k\psi\}_{k\in\mathbb{Z}}$ if and only if $\hat{\phi}\hat{\psi}\chi_\Omega = \chi_\Omega$ a.e. Moreover, if ψ is also such that $|\hat{\psi}| > 0$ a.e. on a connected neighborhood of 0 in $[-\frac{1}{2}, \frac{1}{2})$, and $|\hat{\psi}| = 0$ a.e. otherwise, and $\hat{\phi}\hat{\psi}\chi_\Omega = \chi_\Omega$ a.e. holds, then $\{\pi_k\phi\}_{k\in\mathbb{Z}}$ and $\{\pi_k\psi\}_{k\in\mathbb{Z}}$ are commutative pair of pseudoframe for X .*

3. Generalized Multiresolution Structure

In this section, by applying Theorem 2.3, we are going to construct a generalized multiresolution structure for locally compact abelian groups. First, we define the concept of generalized multiresolution structure (GMS) for $L^2(G)$, where G is a locally compact abelian group.

Let $\pi : \Gamma \rightarrow U(L^2(G))$ be the translation representation and also, let σ be a unitary operator on $L^2(G)$ with the property $\sigma^{-1}\pi_\gamma\sigma = \pi_{\alpha(\gamma)}$, where α is an injective endomorphism on Γ . Also let δ be an automorphism on G such that $\delta(\Gamma) \subset \Gamma$.

Definition 3.1. A generalized multiresolution structure (GMS), $\{V_j, \phi, \psi\}_{j\in\mathbb{Z}}$ of $L^2(G)$ is an increasing sequence of the closed subspaces $V_j \subseteq L^2(G)$ and two elements $\phi, \psi \in L^2(G)$ such that the following conditions hold

1. $\overline{\cup_j V_j} = L^2(G), \cap_j V_j = \{0\}$,
2. $f \in V_j$ if and only if $\sigma f \in V_{j+1}$,
3. $f \in V_0$ implies that $\pi_\gamma f \in V_0$, for all $\gamma \in \Gamma$,
4. $\{\pi_\gamma\phi : \gamma \in \Gamma\}$ is a pseudoframe for V_0 with respect to $\{\pi_\gamma\psi : \gamma \in \Gamma\}$.

Remark 3.2. If $\{\pi_\gamma\phi\}_{\gamma\in\Gamma}$ and $\{\pi_\gamma\psi\}_{\gamma\in\Gamma}$ are dual frames, then GMS is a frame multiresolution analysis. Also, if $\{\pi_\gamma\phi\}_{\gamma\in\Gamma}$ is an exact frame for V_0 and $\psi \in V_0$, then GMS is an multiresolution analysis.

Theorem 3.3. *Suppose that $\{\pi_\gamma\phi\}_{\gamma\in\Gamma}$ is a pseudoframe for V_0 with respect to $\{\pi_\gamma\psi\}_{\gamma\in\Gamma}$ and $V_j := \{f \in L^2(G) : \sigma^{-j}f \in V_0\}$, then $\{\sigma^j\pi_\gamma\phi\}_{\gamma\in\Gamma}$ is a pseudoframe for V_j with respect to $\{\sigma^j\pi_\gamma\psi\}_{\gamma\in\Gamma}$.*

Proof. For $f \in V_j$, we have $\sigma^{-j}f \in V_0$. So $\sigma^{-j}f = \sum_{\gamma\in\Gamma} \langle \sigma^{-j}f, \pi_\gamma\psi \rangle \pi_\gamma\phi$. We have $\sigma^* = \sigma^{-1}$, since σ is unitary. Thus

$$f = \sum_{\gamma\in\Gamma} \langle \sigma^{-j}f, \pi_\gamma\psi \rangle \sigma^j \pi_\gamma\phi = \sum_{\gamma\in\Gamma} \langle f, \sigma^j \pi_\gamma\psi \rangle \sigma^j \pi_\gamma\phi.$$

□

Corollary 3.4. *Let $\phi, \psi \in L^2(G)$ and V_0 has the properties specified in Theorem 2.3 and V_j is similar to Theorem 3.3, then $\{V_j, \phi, \psi\}_j$ forms a GMS for $L^2(G)$.*

Proof. The inclusion $V_j \subseteq V_{j+1}$ follows from the fact that V_j defined by Theorem 3.3 is equivalent to $PW_{\Delta(\delta^j)}$ and $PW_{\Delta} \subseteq PW_{\Delta(\delta)}$. Since $\delta(\Gamma) \subset \Gamma$ we have $\hat{\Gamma}(\delta) \subset \hat{\Gamma}$. Now let $f \notin PW_{\Delta}$. Then $supp \hat{f}$ is not a subset of Δ , so $supp \hat{f}$ is not a subset of $\Delta(\delta)$, consequently, $f \notin PW_{\Delta(\delta)}$. Therefore, $PW_{\Delta} \subseteq PW_{\Delta(\delta)}$.

Trivially the set of all band-limited functions, that their Fourier transform has a bounded support, is dense in $L^2(G)$. On the other hand, the intersection of all band-limited function is the trivial function.

For proving condition 3. of Definition 3.1, note that if $f \in V_0$, then $f = \sum_{\eta \in \Gamma} \langle f, \pi_{\eta} \psi \rangle \pi_{\eta} \phi$. Therefore,

$$\pi_{\gamma} f = \sum_{\eta \in \Gamma} \langle f, \pi_{\eta} \psi \rangle \pi_{\gamma} \pi_{\eta} \phi = \sum_{\eta \in \Gamma} \langle f, \pi_{\eta \gamma^{-1}} \psi \rangle \pi_{\eta} \phi = \sum_{\eta \in \Gamma} \langle \pi_{\gamma} f, \pi_{\eta} \psi \rangle \pi_{\eta} \phi.$$

Thus $\pi_{\gamma} f \in V_0$. Other conditions of Definition 3.1 are valid, obviously. □

Example 3.5. Let $G = \mathbb{R} \times \mathbb{R}$ and $\Gamma = \mathbb{Z} \times \mathbb{Z}$. Let $\phi \in L^2(G)$ be such that

$$\hat{\phi}(\gamma_1, \gamma_2) = \begin{cases} 1 & 0 \leq \gamma_1^2 + \gamma_2^2 \leq \frac{1}{16}, \quad a.e. \\ 2 - 4\sqrt{\gamma_1^2 + \gamma_2^2} & \frac{1}{16} \leq \gamma_1^2 + \gamma_2^2 \leq \frac{1}{4} \quad a.e. \\ 0 & otherwise. \end{cases}$$

Put $\Delta := \{(\gamma_1, \gamma_2) \in \mathbb{R} \times \mathbb{R} : |\hat{\phi}(\gamma_1, \gamma_2)| \geq 1\} = \{(\gamma_1^2, \gamma_2^2) : 0 \leq \gamma_1^2 + \gamma_2^2 \leq \frac{1}{16}\}$ and define $V_0 := PW_{\Delta}$. Now select $\psi \in L^2(G)$ such that

$$\hat{\psi}(\gamma_1, \gamma_2) = \begin{cases} 1 & 0 \leq \gamma_1^2 + \gamma_2^2 \leq \frac{1}{16} \quad a.e. \\ 3 - 8\sqrt{\gamma_1^2 + \gamma_2^2} & \frac{1}{16} \leq \gamma_1^2 + \gamma_2^2 \leq \frac{9}{64} \quad a.e. \\ 0 & otherwise. \end{cases}$$

On Δ , we have $\hat{\phi} \cdot \hat{\psi} = 1$, thus $\{V_j, \phi, \psi\}$ forms a GMS for the space of $L^2(G)$.

Note that for the Haar measure dt on G , and an endomorphism δ , the formation $d(\delta(t))$ induces a Haar measure. Thus there is a positive number $|\delta|$ such that $d(\delta(t)) = |\delta|dt$. It is obvious that $d(\delta^{-1}(t)) = |\delta|^{-1}dt$. Now we define a suitable dilation by $\sigma f(t) := |\delta|^{-\frac{1}{2}} f(\delta(t))$.

Let two complex families $\{h_0(\eta)\}_{\eta \in \Gamma}, \{h_0^*(\eta)\}_{\eta \in \Gamma}, (h_0, h_0^* : \Gamma \rightarrow \mathbb{C})$ are two functions) be such that the following summations are convergent,

$$H_0(\xi) := \sum_{\eta \in \Gamma} h_0(\eta) \overline{\xi(\eta)}, H_0^*(\xi) := \sum_{\eta \in \Gamma} h_0^*(\eta) \overline{\xi(\eta)}, \quad (\xi \in \hat{G}).$$

Proposition 3.6. Suppose H_0 and H_0^* generate ϕ and ψ , respectively, as

$$\phi(t) = \sum_{\gamma \in \Gamma} h_0(\gamma) \sigma \pi_{\gamma} \phi(t), \quad \psi(t) = \sum_{\gamma \in \Gamma} h_0^*(\gamma) \sigma \pi_{\gamma} \phi(t)$$

and $\phi, \psi \in L^2(G)$ have the properties specified in Theorem 2.3. Then $\{\pi_{\gamma} \phi\}_{\gamma \in \Gamma}$ forms a pseudoframe for V_0 with respect to $\{\pi_{\gamma} \psi\}_{\gamma \in \Gamma}$ if and only if

$$H_0.H_0^*\chi_{\Delta(\delta^{-1})} = |\delta|\chi_{\Delta(\delta^{-1})}.$$

Where $\Delta(\delta^{-1}) := \{\gamma\delta^{-1} : \gamma \in \Delta\}$.

Proof. Taking the Fourier transform of ϕ , we get

$$\begin{aligned} \hat{\phi}(\xi) &= \sum_{\gamma \in \Gamma} h_0(\gamma)(\sigma\pi_\gamma\phi)(\xi) \\ &= \sum_{\gamma \in \Gamma} h_0(\gamma) \int_{\hat{G}} |\delta|^{-\frac{1}{2}} \phi(\gamma^{-1}\delta(t))\overline{\xi(t)} dt \\ &= \sum_{\gamma \in \Gamma} h_0(\gamma) \int_{\hat{G}} |\delta|^{-\frac{1}{2}} \phi(t)\overline{\xi(\delta^{-1}(\gamma t))} d(\gamma^{-1}\delta(t)) \\ &= \sum_{\gamma \in \Gamma} h_0(\gamma) \int_{\hat{G}} |\delta|^{-\frac{1}{2}} \phi(t)\overline{\xi(\delta^{-1}(\gamma))\xi(\delta^{-1}(t))} |\delta| dt \\ &= |\delta|^{\frac{1}{2}} H_0(\xi\delta^{-1})\hat{\phi}(\xi\delta^{-1}). \end{aligned}$$

So, $\hat{\phi}(\xi\delta) = |\delta|^{\frac{1}{2}} H_0(\xi)\hat{\phi}(\xi)$. Consequently $H_0.H_0^*\chi_{\Delta(\delta^{-1})} = |\delta|\chi_{\Delta(\delta^{-1})}$. □

4. Affine pseudoframes of $L^2(G)$

We shall denote by W_0 the orthogonal complement of V_0 in V_1 , as usual, in order to split a function f of V_1 into two functions in V_0 and W_0 , respectively.

Definition 4.1. Let $\{V_j, \phi, \psi\}_{j \in \mathbb{Z}}$ be a given GMS and ϕ^*, ψ^* be two functions in $L^2(G)$. We say $\{\pi_\gamma\phi, \pi_\gamma\phi^*\}_{\gamma \in \Gamma}$ is an affine pseudoframe for V_1 with respect to $\{\pi_\gamma\psi, \pi_\gamma\psi^*\}_{\gamma \in \Gamma}$, if and only if

$$f = \sum_{\gamma \in \Gamma} \langle f, \pi_\gamma\psi \rangle \pi_\gamma\phi + \sum_{\gamma \in \Gamma} \langle f, \pi_\gamma\psi^* \rangle \pi_\gamma\phi^*, \quad (f \in V_0).$$

In this case $\{\pi_\gamma\psi, \pi_\gamma\psi^*\}_{\gamma \in \Gamma}$ is called a dual pseudoframe of $\{\pi_\gamma\phi, \pi_\gamma\phi^*\}_{\gamma \in \Gamma}$.

We are going to characterize conditions for which $\{\pi_\gamma\phi, \pi_\gamma\phi^*\}_{\gamma \in \Gamma}$ is an affine pseudoframe for V_1 with respect to $\{\pi_\gamma\psi, \pi_\gamma\psi^*\}_{\gamma \in \Gamma}$. First we have the following.

Proposition 4.2. Let $\{h_1(\gamma)\}_{\gamma \in \Gamma}$ be such that $H_1(\xi) = \sum_{\eta \in \Gamma} h_1(\eta)\overline{\xi(\eta)}$ is convergent, $H_1(0) = 0$ and $H_1 \in L^\infty(\hat{\Gamma})$. Suppose that $\phi \in L^2(G)$ and $\phi(t) = |\delta|^{\frac{1}{2}} \sum_{\gamma \in \Gamma} h_0(\gamma)\phi(\delta(t)\gamma^{-1})$, for the family $\{h_0(\eta)\}_{\eta \in \Gamma}$ such that $H_0(\xi) = \sum_{\eta \in \Gamma} h_0(\eta)\overline{\xi(\eta)}$ is convergent. Then there exists $\psi \in L^2(G)$ such that

$$\psi(t) = |\delta|^{\frac{1}{2}} \sum_{\gamma \in \Gamma} h_1(\gamma)\phi(\delta(t)\gamma^{-1}). \tag{4.1}$$

Proof. Define

$$\hat{\psi}(\omega) = |\delta|^{\frac{1}{2}} H_1(\omega\delta) \prod_{j=2}^{\infty} |\delta|^{\frac{1}{2}} H_0(\omega\delta^{-j}).$$

By the equality $\hat{\phi}(\xi) = H_0(\xi\delta^{-1})\hat{\phi}(\xi\delta^{-1})$, we have

$$\hat{\psi}(\omega) = |\delta|^{\frac{1}{2}}H_1(\omega\delta^{-1})\hat{\phi}(\omega\delta^{-1}) \tag{4.2}$$

Hence $\psi \in L^2(G)$, since $\phi \in L^2(G)$ and $H_1 \in L^\infty(\hat{\Gamma})$. It is now sufficient to use Parsval's Theorem and inverse Fourier transform of (4.2). to obtain (4.1). \square

Suppose A is a subset of \hat{G} . For the next theorem we need the following Γ^\perp -periodic function

$$\Lambda_A(\gamma) = \sum_{\eta \in \Gamma^\perp} \chi_A(\gamma\eta).$$

Theorem 4.3. *Let Δ be the bandwidth of the subspace V_0 defined by Theorem 2.3. The family $\{\pi_\gamma\phi, \pi_\gamma\phi^*\}_{\gamma \in \Gamma}$ forms a pseudoframe for V_1 with respect to $\{\pi_\gamma\psi, \pi_\gamma\psi^*\}_{\gamma \in \Gamma}$ if and only if there are two functions G_0, G_1 in $L^2(\hat{\Gamma})$ such that*

$$\begin{aligned} G_0(\xi)\overline{H_0^*(\xi)}\Lambda_\Delta(\xi) + G_1(\xi)\overline{H_1^*(\xi)}\Lambda_\Delta(\xi) &= 2\Lambda_\Delta(\xi), \text{ a.e.} \\ G_0(\xi)\overline{H_0^*(\xi)1_{\hat{\Gamma}}\delta^{-1}}\Lambda_\Delta(\xi) + G_1(\xi)\overline{H_1^*(\xi)1_{\hat{\Gamma}}\delta^{-1}}\Lambda_\Delta(\xi) &= 0, \text{ a.e.} \end{aligned}$$

Proof. The fact that $\{\psi_{1,\eta}\}_{\eta \in \Gamma}$ generates the elements of V_1 , implies that for any $f \in V_1$,

$$\langle f, \psi_{1,\eta} \rangle = \sum_{\gamma \in \Gamma} \langle f, \pi_\gamma\psi \rangle \langle \pi_\gamma\phi, \psi_{1,\eta} \rangle + \sum_{\gamma \in \Gamma} \langle f, \pi_\gamma\psi^* \rangle \langle \pi_\gamma\phi^*, \psi_{1,\eta} \rangle, \quad (\eta \in \Gamma).$$

Now define

$$\begin{aligned} c_0(\gamma) &= \langle f, \pi_\gamma\psi \rangle, \\ c_1(\gamma) &= \langle f, \psi_{1,\gamma} \rangle, \\ d_0(\gamma) &= \langle f, \pi_\gamma\psi^* \rangle, \end{aligned}$$

and also let

$$g_0(\eta(\delta(\gamma))^{-1}) = \langle \pi_\gamma\phi, \psi_{1,\eta} \rangle, \quad g_1(\eta(\delta(\gamma))^{-1}) = \langle \pi_\gamma\phi^*, \psi_{1,\eta} \rangle.$$

We have

$$c_1(\eta) = \sum_{\gamma \in \Gamma} c_0(\gamma)g_0(\eta(\delta(\gamma))^{-1}) + \sum_{\gamma \in \Gamma} d_0(\gamma)g_1(\eta(\delta(\gamma))^{-1})$$

By taking the Fourier series, we have

$$\begin{aligned}
 C_1(\xi) &= \sum_{\eta \in \Gamma} \sum_{\gamma \in \Gamma} c_0(\gamma) g_0(\eta(\delta(\gamma))^{-1}) \overline{\xi(\eta)} \\
 &+ \sum_{\eta \in \Gamma} \sum_{\gamma \in \Gamma} d_0(\gamma) g_1(\eta(\delta(\gamma))^{-1}) \overline{\xi(\eta)} \\
 &= \sum_{\eta \in \Gamma} \sum_{\gamma \in \Gamma} c_0(\gamma) g_0(\eta) \overline{\xi(\eta \delta(\gamma))} \\
 &+ \sum_{\eta \in \Gamma} \sum_{\gamma \in \Gamma} d_0(\gamma) g_1(\eta) \overline{\xi(\eta \delta(\gamma))} \\
 &= \sum_{\eta \in \Gamma} \sum_{\gamma \in \Gamma} c_0(\gamma) g_0(\eta) \overline{\xi(\eta) \xi(\delta(\gamma))} \\
 &+ \sum_{\eta \in \Gamma} \sum_{\gamma \in \Gamma} d_0(\gamma) g_1(\eta) \overline{\xi(\eta) \xi(\delta(\gamma))} \\
 &= \sum_{\gamma \in \Gamma} c_0(\gamma) \overline{\xi(\delta(\gamma))} \sum_{\eta \in \Gamma} g_0(\eta) \overline{\xi(\eta)} \\
 &+ \sum_{\gamma \in \Gamma} d_0(\gamma) \overline{\xi(\delta(\gamma))} \sum_{\eta \in \Gamma} g_1(\eta) \overline{\xi(\eta)} \\
 &= C_0(\xi \delta) G_0(\xi) + D_0(\xi \delta) G_1(\xi).
 \end{aligned}$$

On the other hand, we know

$$\begin{aligned}
 c_0(\gamma) &= \int_G f(t) \overline{\pi_\gamma \psi(t)} dt = \int_G f(t\gamma) \overline{\psi(t)} dt \\
 &= \int_G f(t\gamma) \sum_{\eta \in \Gamma} |\delta|^{\frac{1}{2}} h_0^*(\eta) \overline{\psi(\eta^{-1} \delta(t))} dt \\
 &= \sum_{\eta \in \Gamma} |\delta|^{\frac{1}{2}} h_0^*(\eta) \int_G f(t) \overline{\psi(\delta(t) \eta^{-1} (\delta(\gamma))^{-1})} dt \\
 &= \sum_{\eta \in \Gamma} |\delta|^{\frac{1}{2}} h_0^*(\eta(\delta(\gamma))^{-1}) \int_G f(t) \overline{\psi(\delta(t) \eta^{-1})} dt \\
 &= \sum_{\eta \in \Gamma} h_0^*(\eta(\delta(\gamma))^{-1}) c_1(\eta).
 \end{aligned}$$

Similarly,

$$d_0(\gamma) = \sum_{\eta \in \Gamma} h_1^*(\eta(\delta(\gamma))^{-1}) c_1(\eta).$$

Now, their Fourier series are

$$\begin{aligned}
 C_0(\xi\delta) &= \sum_{\gamma \in \Gamma} c_0(\gamma) \overline{\xi(\delta(\gamma))} = \sum_{\gamma \in \Gamma} \sum_{\eta \in \Gamma} \overline{h_0^*(\eta(\delta(\gamma))^{-1})} c_1(\eta) \overline{\xi(\delta(\gamma))} \\
 &= \sum_{\eta \in \Gamma} \sum_{\gamma \in \Gamma} c_1(\eta) \overline{h_0^*(\gamma)} \overline{\xi(\eta\gamma^{-1})} \\
 &= \sum_{\eta \in \Gamma} c_1(\eta) \overline{\xi(\eta)} \sum_{\gamma \in \Gamma} \overline{h_0^*(\gamma)} \overline{\xi(\gamma)} \\
 &= \frac{1}{2} \left[\sum_{\eta \in \Gamma} c_1(\eta) \overline{\xi(\eta)} \sum_{\gamma \in \Gamma} \overline{h_0^*(\gamma)} \overline{\xi(\gamma)} \right. \\
 &\quad \left. + \sum_{\eta \in \Gamma} c_1(\eta) \overline{\xi(\eta)} 1_{\Gamma}(\delta^{-1}(\eta)) \sum_{\gamma \in \Gamma} \overline{h_0^*(\gamma)} \overline{\xi(\gamma)} 1_{\Gamma}(\delta^{-1}(\gamma)) \right] \\
 &= \frac{1}{2} [C_1(\xi) \overline{H_0^*(\xi)} + C_1(\xi 1_{\Gamma} \delta^{-1}) \overline{H_0^*(\xi 1_{\Gamma} \delta^{-1})}].
 \end{aligned}$$

Similarly,

$$D_0(\xi\delta) = \frac{1}{2} [C_1(\xi) \overline{H_1^*(\xi)} + C_1(\xi 1_{\Gamma} \delta^{-1}) \overline{H_1^*(\xi 1_{\Gamma} \delta^{-1})}].$$

Combining the above relations, we find that

$$\begin{aligned}
 2C_1(\xi) &= G_0(\xi) [C_1(\xi) \overline{H_0^*(\xi)} + C_1(\xi 1_{\Gamma} \delta^{-1}) \overline{H_0^*(\xi 1_{\Gamma} \delta^{-1})}] \\
 &\quad + G_1(\xi) [C_1(\xi) \overline{H_1^*(\xi)} + C_1(\xi 1_{\Gamma} \delta^{-1}) \overline{H_1^*(\xi 1_{\Gamma} \delta^{-1})}] \\
 &= [G_0(\xi) \overline{H_0^*(\xi)} + G_1(\xi) \overline{H_1^*(\xi)}] C_1(\xi) \\
 &\quad + [G_0(\xi) \overline{H_0^*(\xi 1_{\Gamma} \delta^{-1})} + G_1(\xi) \overline{H_1^*(\xi 1_{\Gamma} \delta^{-1})}] C_1(\xi 1_{\Gamma} \delta^{-1}).
 \end{aligned}$$

Consequently,

$$\begin{aligned}
 G_0(\xi) \overline{H_0^*(\xi)} \Lambda_{\Delta}(\xi) + G_1(\xi) \overline{H_1^*(\xi)} \Lambda_{\Delta}(\xi) &= 2\Lambda_{\Delta}(\xi), \text{ a.e.} \\
 G_0(\xi) \overline{H_0^*(\xi 1_{\Gamma} \delta^{-1})} \Lambda_{\Delta}(\xi) + G_1(\xi) \overline{H_1^*(\xi 1_{\Gamma} \delta^{-1})} \Lambda_{\Delta}(\xi) &= 0, \text{ a.e.}
 \end{aligned}$$

□

Acknowledgments

This research was supported by a grant from Ferdowsi University of Mashhad-Graduate studies No. 29509.

References

- [1] J. J. Benedetto, S. Li, *The theory of multiresolution analyses frames and applications to filter banks*, Appl. Comp. Harm. Anal., **5** (1998), 389-427.

- [2] J. J. Benedetto, S. Li, Multiresolution analysis frames with applications, Proceeding ICASSP'93 Proceedings of IEEE international conference on Acoustics, speech, and signal processing: digital speech processing - Volume III Pages 304-307, 1993
- [3] P. G. Casazza O. Christensen, D. Stoeva, *Frame expansions in separable Banach space*, J. Math. Anal. Appl., **114**(1) (2005), 710–723.
- [4] O. Christensen, *On frame multiresolution analysis*, In "An Introduction to Frames and Riesz Bases", Part of the series Applied and Numerical Harmonic Analysis, 283-311, 2003.
- [5] O. Christensen, *An Introduction to Frames and Riesz Bases*, Birkhäuser, Boston 2003.
- [6] S. Dahlke, *Multiresolution Analysis and Wavelets on Locally Compact Abelian Groups*, Wavelets, Images, and Surface Fitting, P. J. Laurent, A. Le Mehaute, L. L. Schumaker, eds., A. K. Peters, Wellesley, 1994, 141-156.
- [7] I. Daubechies, *Orthonormal bases of compactly supported wavelets*, Commun. Pure Appl. Math., **41**(9) (1988), 909-996.
- [8] I. Daubechies, A. Grossmann, Y. Meyer, *Painless nonorthogonal expansion*, J. Math. Phys. **27** (1986), 1271–1283.
- [9] R. Duffin R, S. Schaeffer, *A class of nonharmonic Fourier series*, Trans. Amer. Math. Soc., **72** (1952), 341–366.
- [10] Yu. A. Farkov, Orthogonal wavelets on locally compact abelian groups, *Funktsional. Anal. i Prilozhen.*, **31**(4) (1997), 86-88; English transl., *Funct. Anal. Appl.*, **31** (1997), 294-296.
- [11] G. B. Folland, *A Course in Abstract Harmonic Analysis*, CRC Press, 1995.
- [12] D. Gabor, *Theory of communications*, J. Inst. Electr. Eng., **93**(26) (1946), 429–457.
- [13] R. A. Kamyabi Gol, R. Raisi Tousi, *The structure of shift invariant spaces on a locally compact abelian group*, J. Math. Anal. Appl., **340**(1) (2008), 219–225.
- [14] R. A. Kamyabi Gol, R. Raisi Tousi, *Some equivalent multiresolution conditions on locally compact abelian groups*, Proc. Math. Sci., **120**(3) (2010), 317–331.
- [15] S. V. Kozyrev, *Wavelet theory as p-adic spectral analysis*, *Izv. Ross. Akad. Nauk Ser. Mat.*, **66**(2) (2002), 149-158. English transl, *Izv. Math.*, **66** (2002), 367–376.
- [16] W. C. Lang, *Wavelet analysis on the Cantor dyadic group*, *Houston J. Math.*, **24**(3)(1998), 533–544.
- [17] S. Li, *The theory of frame multiresolution analysis and its applications*, Ph. D. Thesis, University of Maryland Graduate School, Baltimore, May 1993.
- [18] S. Li, *A theory of generalized multiresolution structure and pseudoframes of translates*, J. Fourier Anal. and Appl. **7**(1) (2001), 23–40.
- [19] S. Li, H. Ogawa, *A theory of pseudoframes for subspaces with applications*, Proc. SPIE 3458, Wavelet Applications in Signal and Imaging Processing VI, 67(1998); doi:10.1117/12.328126.
- [20] S. Li, H. Ogawa, *Pseudoframes for subspaces with applications*, J. Fourier Anal. Appl., **10**(4)(2004), 409–431.
- [21] S. G. Mallat, *Multiresolution approximations and wavelet orthonormal bases of $L^2(\mathbb{R})$* , Trans. Amer. Math. Soc., **315**(1) (1989), 69–87.
- [22] Y. Meyer, *Wavelets and Operators*, Translated by DH Salinger, Cambridge Studies in Advanced Mathematics, 1992.
- [23] D. P. Petersen, D. Middleton, *Sampling reconstruction of wave-number limited functions in N-dimensional Euclidean spaces*, Inf. Control, **5**(4) (1962), 279–323.